# **Nuclear Medicine Preparation Instructions**

#### **Bone Scan**

No preparation, allow 3- 4 hours for the scan (2 part test with minimum 2 hours between parts).

#### **Brain Scan**

No preparation, allow 2 hours for the scan.

#### **Gallium Scan**

No preparation. Injection is on the first day. Scanning (30-90 minutes) will be done 2-3 days post injection.

### **Gastric Emptying Scan**

Fasting for a minimum of 4 hours. Patient must be off certain stomach medication for 48 hours. Please check with Nuclear Medicine Department. Patient will be required to eat egg whites, bread and jam. Test takes approximately 5 hours with breaks.

### **Hepatobiliary Scan**

Fasting for a minimum of 4 hours. Allow 2-3 hours for the scan.

### Liver / Spleen - Sulfur Colloid

No preparation. Allow 1 hour for the scan.

#### Liver / Spleen - Red blood cell

No preparation. Allow 4 hours for the scan (2 part test with 2.5 hours between parts)

### Lung Scan

No preparation. Allow 45 minutes for the scan

#### Parathyroid Scan

No preparation, allow 3-4 hours for the scan

## Renal Scan (with/without diuretic)

Recent Lab work completed (Creatinine level ) is required prior to booking. Patient must be drink a minimum 500 cc of water - a minimum of I hour prior to scan. Patient can void at any time. Allow 45 -90 minutes for scan.

### Salivary Scan

Patient to bring a lemon or lemon juice for the scan. Allow 60 minutes for the scan.

## **Renal Scan with Captopril**

Patient must fast for a minimum of 4 hours prior to exam.

Patient must drink a minimum 500 cc of water and can void at any time.

Patient must be off blood pressure medication for 48 hours and diuretics for 72 hours.

Allow 2 hours for the scan.

Baseline study may be needed one week later.

## **Thyroid Scan/ Uptake**

Patient must discontinue thyroid medication for 1 to 4 weeks.

No CT contrast for up to 8 weeks.

No cough medicine, seaweed or highly iodinated substances for 1 week.

Please speak to nuclear medicine physician if patient is on amiodarone.

Patient has to be able to swallow a capsule for 2 consecutive days.

Allow 45 minutes for the scan.